

On Banach Spaces of Large Density Character and Unconditional Sequences

Eugene Tokarev

Dedicated to the memory of S. Banach.

ABSTRACT. It is shown that any Banach space X of density $\text{dens}(X) \geq \mathfrak{J}_{\omega_1}$ contains an (infinite) unconditional sequence. Here for any ordinal α the cardinal $\mathfrak{J}_\alpha$ is given by induction: $\mathfrak{J}_0 = \aleph_0$; $\mathfrak{J}_{\alpha+1} = \exp(\mathfrak{J}_\alpha)$; $\mathfrak{J}_\alpha = \sup\{\mathfrak{J}_\beta : \beta < \alpha\}$ if α is a limit ordinal, and ω_1 is the first uncountable ordinal.

Every Banach space of density $\text{dens}(X) = \tau \geq \omega_1$ contains an unconditional sequence of cardinality τ if and only if τ is a weakly compact cardinal.

Other results that are concerned large cardinals and unconditional sequences are also presented.

1. Introduction.

The known J. Lindenstrauss' problem [1] was:

Whether every infinite dimensional Banach space contains an (infinite) unconditional sequence?

It was successfully solved (in negative) by W.T. Gowers and B. Maurey [2].

More formerly, J. Ketonen [3] has proved that any Banach space X of dimension $\dim(X) = \varrho$, where ϱ is a Ramsey cardinal (definitions see below) contains a subspace with an unconditional basis of the cardinality ϱ .

Recall that a *dimension* $\dim(X)$ of a Banach space X is the least cardinality of a subset $A \subset X$, which *linear span* $\text{lin}(A)$ is dense in X (equivalently: such $A \subset X$ that a *closure* $\overline{\text{lin}(A)}$, which will be in the future denoted $\text{span}(A)$, is the whole space X). If X is of infinite dimension then its dimension $\dim(X)$ is exactly equal to its *density character* $\text{dens}(X)$ - the least cardinality of a subset $B \subset X$, which is dense in X .

The result of [3] was improved in [4] (see also [5]), where was shown that for any cardinal τ there exists a cardinal $\varkappa(\tau)$ (namely, an P. Erdős' cardinal $\varkappa(\tau) \rightarrow (\tau)_2^{<\omega}$) such that any Banach space of dimension $\geq \varkappa(\tau)$ contains an unconditional sequence of cardinality τ . Cardinals of kind $\varkappa(\tau)$ are too large. It is known that $\iota < \varkappa(\omega) < \varkappa(\omega_1) < \dots < \varkappa(\varkappa) = \varrho$, where ι is the first inaccessible

1991 *Mathematics Subject Classification.* Primary 46B26; Secondary 46B03, 46B04, 46B07, 46B20 03C65, 03C75.

Key words and phrases. Unconditional sequences, Banach spaces, Large cardinals Infinitary logic $\mathfrak{L}_{\omega_1, \omega}$.

cardinal, and ϱ is the first Ramsey cardinal (here and below ω is the first infinite cardinal; ω_1 is the first uncountable cardinal). Moreover, each $\varkappa(\tau)$ is inaccessible itself. So, the existence of $\varkappa(\omega)$, of $\varkappa(\omega_1)$, etc. must be postulated separately.

P. Terenzi [6] asked:

Whether there exists a cardinal τ (without any additional set-theoretical assumptions) such that any Banach space X of dimension $\dim(X) \geq \tau$ contains an infinite unconditional sequence?

By using of some methods from the model theory it may be proved that the existence of $\varkappa(\omega)$ implies that any Banach space of dimension $\geq \aleph_{\omega_1}$ contains an unconditional sequence (the cardinal $\aleph_{\omega_1}$, which existence does not depend on any set-theoretical hypotheses, will be defined later).

Below it will be shown that, independently of the existence (or non-existence) of $\varkappa(\omega)$, any Banach space X of $\dim(X) \geq \aleph_{\omega_1}$ contains an unconditional sequence.

A sequence $\{x_\alpha : \alpha < \tau\}$ of elements of a Banach space X will be called a τ -sequence. The following question is of a separate interest:

What cardinals τ has the property: each Banach space X of dimension τ contains an unconditional τ -sequence?

It will be shown that a cardinal τ has a connected property (namely, each Banach space of dimension τ contains an subsymmetric τ -sequence) if and only if τ is a weakly compact cardinal. Hence, the same property of Ramsey cardinals (cf. [1]) is a consequence of their weak compactness.

By the way, some partial results in the S. Banach's problem on the separable quotient will be obtained.

2. Definitions and notations

Ordinals will be denoted by small Greece letters α, β, γ . *Cardinals* are identified with the least ordinals of a given cardinality and are denoted either $\iota, \tau, \varkappa, \varrho, \sigma$, or by using Hebrew letters $\aleph, \beth$ (may be with indices). As usual, ω and ω_1 denote respectively the first infinite and the first uncountable cardinals (=ordinals).

For a cardinal τ its *predecessor* (i.e. the least cardinal that is strongly greater than τ) is denoted by τ^+ .

The *cofinality* of τ , $cf(\tau)$ is the least cardinality of a set $A \subset \tau$ such that $\tau = \sup A$.

Let A, B be sets. A symbol ${}^B A$ denotes a set of all functions from B to A .

In a general case, a cardinality of the set ${}^B A$ is denoted either $\text{card}(A)^{\text{card}(B)}$ or $\varkappa^\tau$, if $\text{card}(A) = \varkappa$; $\text{card}(B) = \tau$.

A symbol $\exp(\tau)$ (or, equivalently, 2^τ) denotes a cardinality of a set ${}^\tau 2$ of all subsets of τ .

Let J be a set; $n < \omega$. Let

$$[J]^n = \{A \subset J : \text{card}(A) = n\}$$

J may be considered as a well-ordered set (say, by a relation ' $<$ '). So, $[J]^n$ may be regarded as a set of all n -tuples $a_0 < a_1 < \dots < a_{n-1}$ where $a_i \in J$ ($i < n$).

For a cardinal σ a function $P : [J]^n \rightarrow \sigma$ is called a *partition* of $[J]^n$ (on σ parts).

If $P : [J]^n \rightarrow \sigma$; $H \subset [J]^n$ and the function P is a constant on $[H]^n$, i.e., if all elements of $[H]^n$ belong to the same class of the partition P , then the set H is said to be *homogeneous for P* .

The P. Erdős' denotation

$$\kappa \rightarrow (\tau)_\sigma^n$$

will be used as a shortness for the following assertion:

For every partition $P : [\kappa]^n \rightarrow \sigma$ there exists a homogeneous for P subset $H \subset \kappa$ of cardinality τ .

The another Erdős' denotation

$$\kappa \rightarrow (\tau)_\sigma^{<\omega}$$

means:

For any family $\{P_n : n < \omega\}$ of partitions $P_n : [\kappa]^n \rightarrow \sigma$ there exists a set $H \subset \kappa$, which is homogeneous for all of P_n 's.

Let τ be a cardinal. An *Erdős' cardinal* $\kappa(\tau)$ is the least cardinal κ such that

$$\kappa \rightarrow (\tau)_2^{<\omega}.$$

A cardinal σ is said to be *weakly compact* if

$$\sigma \rightarrow (\sigma)_2^2.$$

A cardinal ρ is said to be a *Ramsey cardinal* if

$$\rho \rightarrow (\rho)_2^{<\omega}.$$

A cardinal v is said to be *measurable* if there exists a countably additive function (measure) $\mu : v \rightarrow \{0, 1\}$, defined on all subsets of v , such that

$$\mu(v) = 1; \mu(A) = 0 \text{ for every subset } A \subset v \text{ of cardinality } \text{card}(A) < v.$$

Any Ramsey cardinal is weakly compact. Any measurable cardinal is weakly compact and is a Ramsey cardinal.

Let α be an ordinal; τ be a cardinal. A cardinal $\mathfrak{J}_\alpha(\tau)$ is given by the induction:

$$\mathfrak{J}_0(\tau) = \tau; \quad \mathfrak{J}_{\alpha+1}(\tau) = \exp(\mathfrak{J}_\alpha(\tau));$$

$$\mathfrak{J}_\alpha(\tau) = \sup\{\mathfrak{J}_\beta(\tau) : \beta < \alpha\},$$

if α is a *limit ordinal* (i.e. if α is not of kind $\alpha = \beta + 1$ for some ordinal β).

In a case $\tau = \omega$ ($= \aleph_0$) it will be written $\mathfrak{J}_\alpha$ instead of $\mathfrak{J}_\alpha(\omega)$.

Recall that a cardinal $\aleph_\alpha$ is given by

$$\aleph_0 = \omega; \aleph_{\alpha+1} = (\aleph_\alpha)^+; \aleph_\alpha = \sup\{\aleph_\beta : \beta < \alpha\}, \text{ if } \alpha \text{ is a limit ordinal.}$$

The *general continuum hypothesis* (GCH) is a conjecture:

$$\aleph_\alpha = \mathfrak{J}_\alpha \text{ for every ordinal } \alpha.$$

The *continuum hypotheses* (CH) asserts that $\aleph_1 = \mathfrak{J}_1$.

In the future the following Erdős-P. Rado theorem [7] will be used:

For every infinite cardinal κ

$$(\mathfrak{J}_n(\kappa))^+ \rightarrow (\kappa^+)_\kappa^{n+1}.$$

Let $\mathcal{B}$ be a (proper) class of all Banach spaces. Let $X \in \mathcal{B}$; $\dim(X) = \kappa$; α be an infinite limit ordinal (so, α may be considered as a cardinal); $\alpha \leq \kappa$. An α -sequence $\{x_\beta : \beta < \alpha\}$ of elements of X is said to be

- *Almost spreading*, if for any $n < \omega$, any $\varepsilon > 0$, any scalars $\{a_k : k < n\}$ and any choosing of $i_0 < i_1 < \dots < i_{n-1} < \alpha$; $j_0 < j_1 < \dots < j_{n-1} < \alpha$ the following inequality holds:

$$(1 - \varepsilon) \left\| \sum_{k < n} a_k x_{j_k} \right\| \leq \left\| \sum_{k < n} a_k x_{i_k} \right\| \leq (1 + \varepsilon) \left\| \sum_{k < n} a_k x_{j_k} \right\|;$$

- *Spreading*, if the previous is remain valid for $\varepsilon = 0$;
- *C-unconditional*, where $C < \infty$ is a constant, if

$$C^{-1} \left\| \sum_{k < n} a_k \epsilon_k x_{i_k} \right\| \leq \left\| \sum_{k < n} a_k x_{i_k} \right\| \leq C \left\| \sum_{k < n} a_k \epsilon_k x_{i_k} \right\|$$

for any choosing of $n < \omega$; $\{a_k : k < n\}$; $i_0 < i_1 < \dots < i_{n-1} < \alpha$ and of signs $\{\epsilon_k \in \{+, -\} : k < n\}$.

- *Unconditional*, if it is *C-unconditional* for some $C < \infty$.
- *Symmetric*, if for any $n < \omega$, any finite subset $I \subset \alpha$ of cardinality n , any rearrangement ς of elements of I and any scalars $\{a_i : i \in I\}$,

$$\left\| \sum_{i \in I} a_i z_i \right\| = \left\| \sum_{i \in I} a_{\varsigma(i)} z_i \right\|.$$

- *Subsymmetric*, if it is both spreading and 1-unconditional.

Let $C < \infty$ be a constant. Two α -sequences $\{x_\beta : \beta < \alpha\}$ and $\{y_\beta : \beta < \alpha\}$ are said to be *C-equivalent* if for any finite subset $I = \{i_0 < i_1 < \dots < i_{n-1}\}$ of α and for any choosing of scalars $\{a_k : k < n\}$

$$C^{-1} \left\| \sum_{k < n} a_k x_{i_k} \right\| \leq \left\| \sum_{k < n} a_k y_{i_k} \right\| \leq C \left\| \sum_{k < n} a_k x_{i_k} \right\|.$$

Two α -sequences $\{x_\beta : \beta < \alpha\}$ and $\{y_\beta : \beta < \alpha\}$ are said to be equivalent if they are *C-equivalent* for some $C < \infty$.

It is known (cf. [8]) that for any spreading sequence $\{x_n : n < \omega\}$ the sequence of differences $\{x_{2m+1} - x_{2m} : m < \omega\}$ is 1-unconditional (and, hence, the same is true for any α -sequence when $2m+1$ and $2m$ are replaced with corresponding ordinals). Clearly, if $X \in \mathcal{B}$ contains an almost spreading α -sequence then X contains an *almost unconditional* (i.e. *C-unconditional* for every $C > 1$) α -sequence.

REMARK 1. *The same definitions may be used in a case when instead of α -sequences will be regarded families $\{x_i : i \in I\} \subset X$ indexed by elements of a linearly ordered set $\langle I, \ll \rangle$. In a such case it will be said about spreading families, unconditional families and so on.*

3. On subspaces with unconditional bases in Banach spaces of large dimension

Let $X \in \mathcal{B}$; $\dim(X) = \mathfrak{J}_{\omega_1}$; $\{x_\alpha : \alpha < \mathfrak{J}_{\omega_1}\}$ be a sequence of linearly independent elements of X of norm $\|x_\alpha\| = 1$ for all α , such that $X = \text{span}\{x_\alpha : \alpha < \mathfrak{J}_{\omega_1}\}$. Let $\text{span}\{x_\beta : \beta < \gamma\} = X_\gamma$ for all $\gamma < \mathfrak{J}_{\omega_1}$. It will be shown that X contains an almost spreading sequence (i.e., ω -sequence). This would imply that X contains an unconditional sequence.

THEOREM 1. *Any Banach space of dimension $\mathfrak{J}_{\omega_1}$ contains an infinite dimensional subspace with an unconditional basis.*

PROOF. Let $\dim(X) = \mathfrak{J}_{\omega_1}$; $E_{\omega_1} = \{x_\alpha : \alpha < \mathfrak{J}_{\omega_1}\}$ - a sequence of linearly independent elements of X of norm $\|x_\alpha\| = 1$ for all α such that

$$X = \text{span}\{x_\alpha : \alpha < \mathfrak{J}_{\omega_1}\}$$

Let $\gamma < \omega_1$. Denote

$$E_\gamma = \{x_\alpha : \alpha < \mathfrak{I}_\gamma\}; \quad X_\gamma = \text{span}\{x_\beta : \beta < \mathfrak{I}_\gamma\}.$$

Let $\Psi = \{\psi_N : N < \omega\}$ be a numeration of all finite sequences of rational numbers (i.e., of elements of $\cup\{\mathbb{Q}^n : n < \omega\}$).

For $\psi = \{a_0, a_1, \dots, a_n\} \in \Psi$ let $lh(\psi) = n = \text{card}(\psi) - 1$

It will be convenient to underline the correspondence between a number N of a given $\psi_N \in \Psi$ in a mentioned numeration and its length $lh(\psi)$ in a following way.

Let

$$\psi_N = \{a_0^{(N)}, a_1^{(N)}, \dots, a_{n_N}^{(N)}, r^{(N)}\}.$$

To each $\psi_N \in \Psi$ will be assigned two formal symbols: ϕ_N^+ and ϕ_N^- .

It will be said that a sequence $\mathfrak{x} = \{x_{i_0}, x_{i_1}, \dots, x_{i_N}\} \subset E_{\omega_1}$, where $i_0 < i_1 < \dots < i_N$, satisfies ϕ_N^+ , symbolically: $\mathfrak{x} \models \phi_N^+$ (resp., ϕ_N^- ; in symbols: $\mathfrak{x} \models \phi_N^-$) if

$$\left\| \sum_{k < n_N} a_k^{(N)} x_{i_k} \right\| \geq r^{(N)}$$

(respectively, if

$$\left\| \sum_{k < n_N} a_k^{(N)} x_{i_k} \right\| < r^{(N)}).$$

For any subset $G \subset E_{\omega_1}$ it will be written $G \models \phi_N^+$ (resp., $G \models \phi_N^-$) if $\mathfrak{x} \models \phi_N^+$ (resp., $\mathfrak{x} \models \phi_N^-$) for all ordered n_N -tuples $\mathfrak{x} = \{x_{i_0}, x_{i_1}, \dots, x_{i_N}\}$ from G .

Let us show by an induction that following is true:

- *There exists a sequence of signs $\{\epsilon_k \in \{+, -\} : k < n\}$, a confinal subset $F_n \subset \omega_1$ and a sequence of subsets $G_\alpha \subset E_\alpha$, $\alpha \in F_n$ such that for any $\alpha \in F_n$*
 1. $\text{card}(G_\alpha) \geq \mathfrak{I}_\gamma$, where α is γ 's element of F_n ;
 2. $G_\alpha \models \phi_m^{\epsilon_m}$ for all $m < n$.

This assertion is obviously valid for $n = 0$, $F_1 = \omega_1$ and $G_\alpha = E_\alpha$.

Assume that $\epsilon_0, \epsilon_1, \dots, \epsilon_{n-1}$, F_n , G_α and $\alpha \in F_n$ are already chosen. Let ψ_n be of length $lh(\psi_n) = p_n \geq 1$.

Let $F'_n \subset F_n$ consists of all elements of F_n that are of kind $\beta + (p_n - 1)$ for some ordinal β :

$$F'_n = \{\alpha \in F_n : \text{there exists } \beta \in F_n \text{ such that } (\alpha \setminus \beta) \cap F_n \text{ consists exactly of } p_n - 1 \text{ elements}\}.$$

Clearly, F'_n is confinal in ω_1 . Let α be a γ 's element of F'_n . Then $\text{card}(G_\alpha) > \mathfrak{I}_{\gamma + p_n - 1}$.

The given ψ_n generates a partition of $[G_\alpha]^{p_n}$ in two parts, G_+ and G_- in dependence of $(x_{i_k})_{k < p_n} \models \phi_n^\epsilon$, where $\epsilon \in \{0, 1\}$. Hence, according to the Erdős-Rado theorem, there exists a subset $H_\alpha \subset G_\alpha$, which is homogeneous for this partition and has a cardinality $\text{card}(H_\alpha) = \mathfrak{I}_\gamma$. Clearly, this is true for every $\alpha \in F_n$. Since there exists only two possibilities for $\epsilon \in \{+, -\}$, there exists a confinal subset $F_{n+1} \subset F_n$ such that for all its elements $\gamma \in F_{n+1}$ all p_m -tuples $\{x_{i_0}, x_{i_1}, \dots, x_{i_{p_m}}\}$ ($m \leq n + 1$) satisfy the same condition of kind $\phi_m^{\epsilon_m}$.

So, we choose ϵ , F_{n+1} and $\{H_\alpha : \alpha \in F_{n+1}\}$. This complete the induction.

As a result, it will be chosen a sequence of signs $\{\epsilon_k \in \{+, -\} : k < \omega\}$, which, together with the given order on $\cup\{\mathbb{Q}^n : n < \omega\}$ and the norm of X defines in a unique way (up to an almost isometry) a subspace of X with an approximately

spreading basis (x_n) (for which may be chosen any subset of $\cap F_n$). Of course, this intersection is non empty because of $\text{card}(F_n) = \omega_1$ for all $n < \omega$ and since $\text{cf}(\omega_1) \neq \omega$. $\square$

REMARK 2. *In an analogous way it may be proved that any Banach space of dimension $\geq \mathfrak{J}_{\tau+}$ contains an unconditional τ -sequence.*

It is conceivable that any Banach space of inaccessible dimension contains an unconditional sequence of inaccessible cardinality.

The validity of this assertion is not known. A weaker version will be stated below. Namely, it will be shown that if σ is the first weakly compact cardinal then any Banach space of dimension $\geq \sigma$ contains an unconditional σ -sequence.

4. On separable quotient spaces of Banach spaces

There is a long standing Banach's problem:

Whether every infinite dimensional Banach space X has a separable quotient space?

It easy to point out a class $\mathcal{K}$ of cardinals such that any Banach space of dimension $\varkappa \in \mathcal{K}$ has a separable quotient.

Let $X \in \mathcal{B}$; $\dim(X) = \varkappa$; $\{x_\alpha : \|x_\alpha\| = 1; \alpha < \varkappa\}$ - a $\varkappa$ -sequence of linearly independent elements of X ; $\text{span}\{x_\alpha : \alpha < \varkappa\} = X$. Let $\text{span}\{x_\beta : \beta < \gamma\} = X_\gamma$ for all $\gamma < \varkappa$.

Certainly, $X_\alpha \hookrightarrow X_\gamma$ (recall that a symbol $Y \hookrightarrow Z$ denotes that Y is a subspace of Z) for any $\alpha < \beta < \varkappa$. Obviously, X may be represented as the closure of the union of the chain $X_0 \hookrightarrow X_1 \hookrightarrow \dots \hookrightarrow X_\alpha \hookrightarrow X_\gamma \hookrightarrow \dots$:

$$X = \overline{\cup\{X_\alpha : \alpha < \varkappa\}}.$$

THEOREM 2. *If a cardinal $\varkappa$ is of countable cofinality, $\text{cf}(\varkappa) = \omega$, then every Banach space of dimension $\varkappa$ has a separable quotient space.*

PROOF. Let $\text{cf}(\varkappa) = \omega$. Then there exists a countable sequence $\varkappa_0 < \varkappa_1 < \dots < \varkappa_n < \dots < \varkappa$ such that $\sup\{\varkappa_n : n < \omega\} = \varkappa$. Hence X may be represented as the closure of the union of a countable chain $X_{\varkappa_0} \hookrightarrow \dots \hookrightarrow X_{\varkappa_n} \hookrightarrow$:

$$X = \overline{\cup\{X_{\varkappa_n} : n < \omega\}}.$$

According to [9], any Banach space that has a such representation has also a separable quotient space. $\square$

THEOREM 3. *Any B -convex Banach space W of dimension $\geq \mathfrak{J}_{\omega_1}$ has a separable quotient space.*

PROOF. Let $\dim(W) \geq \mathfrak{J}_{\omega_1}$. Then its conjugate also is of dimension $\dim(W^*) \geq \mathfrak{J}_{\omega_1}$ and thus contains an unconditional sequence, which spans in W^* either a reflexive subspace, or a subspace isomorphic to l_1 or a subspace which is isomorphic to c_0 . Let $Z \hookrightarrow W^*$ be reflexive. Every its separable subspace $Z_0 \hookrightarrow Z$ is reflexive too. Since Z_0 is weakly* closed, clearly, W has a quotient that is isometric to $(Z_0)^*$.

If W^* contains a subspace isomorphic to c_0 then (since W^* is a conjugate space) W^* contains a subspace isomorphic to $L_1[0, 1]$, which contains reflexive subspaces. Let W^* contains a subspace Y isomorphic to l_1 . Consider its weak* closure

$((Y)_\perp)^\perp$, where for a subset $A \subset W^*$ a symbol $(A)_\perp$ denotes its lower annihilator

$$(A)_\perp = \{w \in W : a(w) = 0 \text{ for all } a \in A\};$$

and for a subset $B \subset W$ a symbol $(B)^\perp$ denotes its upper annihilator

$$(B)^\perp = \{w^* \in W^* : w^*(b) = 0 \text{ for all } b \in B\}.$$

Clearly, $((Y)_\perp)^\perp$ is an $L_1(\mu)$ -space. If a measure μ is discrete then W^* contains a weakly* closed subspace, isomorphic to l_1 and, of course, W has a separable quotient. If a measure μ contains a nontrivial dispersed part, then W^* contains a subspace isomorphic to $L_1[0, 1]$ and, as was mentioned before, has a separable quotient too. $\square$

5. Banach spaces of weakly compact dimension

Recall that if a cardinal $\varkappa$ is weakly compact, i.e. if $\varkappa \rightarrow (\varkappa)_2^2$, then $\varkappa \rightarrow (\varkappa)_\sigma^n$ for all $n < \omega$ and all $\sigma < \varkappa$. Any weakly compact cardinal $\varkappa$ is larger than the first inaccessible one, moreover, $\varkappa$ is inaccessible itself and is the $\varkappa$'s inaccessible cardinal.

Hence, $\varkappa \rightarrow (\varkappa)_{\mathfrak{I}_\alpha}^n$ for each $n < \omega$ and $\alpha < \iota$, where ι is the first inaccessible cardinal.

THEOREM 4. *Let $\varkappa$ be a weakly compact cardinal; $X \in \mathcal{B}$ and $\dim(X) = \varkappa$. Then X contains an unconditional $\varkappa$ -sequence.*

PROOF. Let $\{x_\gamma : \gamma < \varkappa\}$ be a sequence of linearly independent elements of X that spans X . Fix $n < \omega$. Let $\{a_0, \dots, a_{n-1}\}$ are scalars, $b \in \mathbb{R}^+$ - a positive real number. There exists a continuum number of choosing of $\phi = \{n, a_0, \dots, a_{n-1}\}$. The set Φ of such elements ϕ may assumed to be well-ordered; let α_ϕ be the ordinal number of ϕ in this ordering.

It will be said that a finite sequence $\gamma_0 < \gamma_1 < \dots < \gamma_n < \varkappa$ belongs to a class C_α^b if

$$\left\| \sum_{k=0}^n a_k x_{\gamma_k} \right\| = b.$$

For a fixed α a set of all classes C_α^b forms a partition of $[\varkappa]^n$ on continuum parts. Since $\varkappa$ is weakly compact, there exists a homogeneous subset $K(\alpha) \subset \varkappa$ for this partition; $\text{card}(K(\alpha)) = \varkappa$.

Next we proceed by the induction. Assume that sets $\{K(\beta) : \beta < \alpha\}$ are already chosen. If $\alpha = \gamma + 1$ for some γ then $K(\alpha)$ is defined to be a homogeneous subset of $K(\gamma)$ with respect to the partition $\{C_{\gamma+1}^b : b \in \mathbb{R}^+\}$. If α is a limit ordinal then $K(\alpha) = \cap \{K(\beta) : \beta < \alpha\}$. Each $K(\alpha)$ is of cardinality $\varkappa$. Clearly, $\cap \{K(\beta) : \beta < \varkappa\}$ is of cardinality $\varkappa$ too, is well ordered and, hence, is a desired spreading $\varkappa$ -sequence. $\square$

The converse result is also true.

THEOREM 5. *For every cardinal $\varkappa$, which is not weakly compact, there exists a Banach space X of dimension $\dim(X) = \varkappa$, which does not contain any unconditional $\varkappa$ -sequence.*

PROOF. Let W be an abstract set of cardinality κ . On W may be defined a linear order, say, $\ll$, such that $\langle W, \ll \rangle$ contains neither strictly increasing nor strictly decreasing κ -sequences with respect to this order. Let X_0 be a vector space of all formal finite linear combinations of elements of W with real scalars. Elements of X_0 are formal sums $x = \sum_{i \in I} a_i w_i$, where I is finite. Let $\{y_n : n < \omega\}$ be a spreading sequence of elements of some Banach space Y . It will be assumed that $\{y_n : n < \omega\}$ is not equivalent to any symmetric sequence.

For any element $x = \sum_{i \in I} a_i w_i$ let $\|x\| = \|\sum_{i \in I} a_i z_i\|_Y$. Let X be a completion of X_0 with respect to this norm. Of course, $\dim(X) = \kappa$. Clearly, $\{w_w : w \in W\}$ is a spreading family, indexed by itself. If $\{s_\alpha : \alpha < \kappa\}$ is a spreading κ -sequence of elements of X , then it must be equivalent to a some block-sequence with respect to $\{w_w : w \in W\}$. However this is impossible for our choosing of W . $\square$

This result may be improved. However, a power of the improvement essentially depends on a model of the set theory, which lies in a base of the whole Banach space theory.

THEOREM 6. *For any cardinal $\tau \geq \omega$ there exists a Banach space X of dimension $\dim(X) = 2^\tau$ ($= \beth_1(\tau)$) that does not contain any subsymmetric τ^+ -sequences.*

PROOF. A "Banach" part of the theorem is similar to the previous one. A set-theoretical part is to find on a set I of a cardinality 2^τ a such linear order, say $\ll$, that $\langle I, \ll \rangle$ contains neither strictly increasing nor strictly decreasing τ^+ -sequences. If $\tau = \omega$, for $\ll$ may be chosen the natural order on the set $\mathbb{R}$ of real numbers; if $\tau > \omega$, as $\ll$ may be regarded a lexicographical ordering on a set ${}^\tau 2$ (of all functions from τ to a set $\{0, 1\}$). $\square$

6. A classification of Banach spaces

Preceding results shows that it is of sense to introduce a partition of the class $\mathcal{B}$ of all Banach spaces by sets

$$\mathcal{B}_\kappa = \{Y \in \mathcal{B} : \dim(Y) = \kappa\},$$

which may be regarded as an analogue of horizontal strips of some generalized coordinate system for Banach spaces. Below were presented results, which demonstrate dependence of properties of Banach spaces on their dimension.

As vertical strips of this generalized coordinate system may be regarded classes of finite equivalence.

DEFINITION 1. *Let X, Y be Banach spaces. X is said to be finitely representable in Y , shortly: $X <_f Y$, if for every $\varepsilon > 0$ and for every finite dimensional subspace A of X there is a subspace B of Y and an isomorphism $u : A \rightarrow B$ such that $\|u\| \|u^{-1}\| < 1 + \varepsilon$. X and Y are said to be finitely equivalent, $X \sim_f Y$, if $X <_f Y$ and $Y <_f X$. So, every Banach space X generates a class of finite equivalence*

$$X^f = \{Y \in \mathcal{B} : X \sim_f Y\}.$$

It will be shown that for any infinite dimensional Banach space X the corresponding class X^f is proper: it contains spaces of arbitrary large dimension.

For a proof there will be used ultrapower of Banach spaces.

DEFINITION 2. Let I be a set; $\text{Pow}(I)$ be a set of all its subsets. An ultrafilter D over I is a subset of $\text{Pow}(I)$ with following properties:

- $I \in D$;
- If $A \in D$ and $A \subset B \subset I$, then $B \in D$;
- If $A, B \in D$ then $A \cap B \in D$;
- If $A \in D$, then $I \setminus A \notin D$.

DEFINITION 3. Let I be a set; D be an ultrafilter over I ; $\{X_i : i \in I\}$ be a family of Banach spaces. An ultraproduct $(X_i)_D$ is given by a quotient space

$$(X)_D = l_\infty(X_i, I) / N(X_i, D),$$

where $l_\infty(X_i, I)$ is a Banach space of all families $\mathfrak{x} = \{x_i \in X_i : i \in I\}$, for which

$$\|\mathfrak{x}\| = \sup\{\|x_i\|_{X_i} : i \in I\} < \infty;$$

$N(X_i, D)$ is a subspace of $l_\infty(X_i, I)$, which consists of such $\mathfrak{x}$'s that

$$\lim_D \|x_i\|_{X_i} = 0.$$

If all X_i 's are all equal to a space $X \in \mathcal{B}$ then an ultraproduct is said to be an *ultrapower* and is denoted by $(X)_D$.

An operator $d_X : X \rightarrow (X)_D$ that asserts to any $x \in X$ an element $(x)_D \in (X)_D$, which is generated by a stationary family $\{x_i = x : i \in I\}$, is called the *canonical embedding* of X into its ultrapower $(X)_D$.

It is well-known that a Banach space X is finitely representable in a Banach space Y if and only if there exists such ultrafilter D (over $I = \cup D$) that X is isometric to a subspace of the ultrapower $(Y)_D$ (cf. [9], an excellent exposition on ultrapowers in the Banach space theory. Below will be presented some important results on ultrapowers that does not contained in [9]).

DEFINITION 4. Let D be an ultrafilter, κ be an cardinal. An ultrafilter D is said to: be

- κ -regular if there exists a subset $G \subset D$ of cardinality $\text{card}(G) = \kappa$ such that any $i \in I$ belongs only to a finite number of sets $e \in G$;
- κ -complete if an intersection of a nonempty subset $G \subset D$ of cardinality $< \kappa$ belongs to D , i.e., if

$$G \subset D \text{ and } \text{card}(G) < \kappa \text{ implies that } \bigcap G \in D;$$

- Principal (or non-free), if there exists $i \in I$ such that $D = \{e \subset I : i \in e\}$ (such ultrafilter is said to be generated by $i \in I$);
- Free, if it is not a principal one;
- Countably incomplete if it is not ω_1 -complete.

THEOREM 7. Let X be a Banach space, $\dim(X) = \kappa$; D be an ultrafilter. The canonical embedding $d_X : X \rightarrow (X)_D$ maps X **on** $(X)_D$ if and only if D is κ^+ -complete.

PROOF. Let D be κ^+ -complete. If $\text{card}(I) \leq \kappa$ then D is non-free and, hence, $d_X X = (X)_D$. Assume that $\text{card}(I) > \kappa$ and that $(x_i)_D \in (X)_D \setminus d_X X$. Of course, $(x_i)_D$ is generated by a such family $(x_i)_{i \in I}$ that for any $\varepsilon > 0$

$$I_0(\varepsilon) = \{i \in I : \|x_i - x\| > \varepsilon\} \in D$$

for any $x \in X$. Consider a function $F : I_0 \rightarrow X$, which is given by $f(i) = x_i$. Since $\dim(X) = \kappa$, a partition

$$I = \cup \{f^{-1}(x) : x \in I_0(\varepsilon)\} \cup \{I \setminus I_0(\varepsilon)\}$$

participate I on τ sets where $\tau < \kappa^+$. Since D is κ^+ -complete, one of sets of the partition belongs to D . Because of $I_0(\varepsilon) \in D$, $I \setminus I_0(\varepsilon) \notin D$. Hence there exists such $x \in X$ that $f^{-1}(x) \in D$. Clearly, this contradicts with the assumption $(x_i)_D \in (X)_D \setminus d_X X$.

Conversely, assume that $d_X X = (X)_D$. Since $\dim(X) = \kappa$ there exists $\varepsilon > 0$ and a set $A \subset X$ such that $\text{card}(A) = \kappa$ and $\|a - b\| \geq \varepsilon$ for all $a \neq b \in A$.

Let $I = \cup \{I_a : a \in A\}$ be a partition of I on $\tau = \text{card}(A) < \kappa^+$ parts. Let a function $F : I \rightarrow A$ is given by:

$$F(i) = a \text{ if and only if } i \in I_a.$$

Then $(F(i))_D \in (X)_D = d_X X$, and, hence, $(F(i))_D = d_X(a)$ for some $a \in A$. Because of $\|a - b\| \geq \varepsilon$ for all $a \neq b \in A$, $F^{-1}(a) \in D$. However, by our assumption, $F^{-1}(a) = I_a$. Thus, $I_a \in D$ and, since a partition of I by τ parts was arbitrary, D is κ^+ -complete. $\square$

COROLLARY 1. *Let X be a Banach space, D be an ultrafilter. then*

1. *If $\dim(X) < \omega$ then $(X)_D = X$;*
2. *If $\omega \leq \dim(X) < \nu$ where ν is the first measurable cardinal, and D is a free ultrafilter then $(X)_D \setminus d_X X \neq \emptyset$;*
3. *If X is of infinite dimension and ultrafilter D is countably incomplete, then $(X)_D \setminus d_X X \neq \emptyset$ too.*

PROOF. Since any ultrafilter is ω -complete (by definitions), all results follows from the preceding theorem. $\square$

Nevertheless, by choosing an ultrafilter, a difference $(X)_D \setminus d_X X$ may be maiden arbitrary large.

THEOREM 8. *Let $X \in \mathcal{B}_\kappa$; D be a τ -regular ultrafilter over a set I of cardinality $\text{card}(I) = \tau$. Then*

$$\dim((X)_D) = (\dim(X))^\tau = \kappa^\tau.$$

PROOF. Let $A' \subset X$ be a set of cardinality κ which is dense in X . Immediately,

$$\dim((X)_D) \leq \dim(l_\infty(X, I)) \leq \text{card}({}^I A) = (\text{card}(A))^{\text{card}(I)} = \kappa^\tau.$$

Assume now that $A \subset A'$ is a set of the same cardinality κ such that for some $\varepsilon > 0$ and any $a, b \in A$, $a \neq b$, $\|a - b\| \geq \varepsilon$. Let $\{a_i : i < \kappa\}$ be a numeration of elements of A . Any finite subset $s = \{x_0, \dots, x_{n-1}\} \subset A$ spans a finite dimensional subspace X_s of X . A set B of all subspaces of kind X_s of X is also of cardinality κ . Let $\{X_\alpha : \alpha < \kappa\}$ be a numeration of B . Consider an ultraproduct $(X_\alpha)_D$. Since $X_\alpha \hookrightarrow X$ for all $\alpha < \kappa$, it may be assumed that $(X_\alpha)_D \hookrightarrow (X)_D$ and, hence, $\dim((X_\alpha)_D) \leq \dim((X)_D)$. Thus, to prove the theorem it is enough to show that $\dim((X_\alpha)_D) \geq \kappa^\tau$.

From τ -regularity of D follows that there exists a subset $G \subset D$, $\text{card}(G) = \tau$, such that any $i \in I$ belongs only to a finite numbers of sets $e \in D$. Define on G a

linear order, say, $\ll$. Let $g : G \rightarrow A$ be a function. Let a function $f_g : I \rightarrow B$ be given by

$$f_g(i) = \text{span}\{a_{g(e_k(i))} : k < n_i\} = B_i,$$

where n_i is a cardinality of a set of those sets $e \in G$, to which i belongs; $\{e_0(i) \ll \dots \ll e_{n_i}(i)\}$ is a list of them. Let $h : G \rightarrow A$ be any other function: $h \neq g$. Then there exists $e \in G$ such that $h(e) \neq g(e)$. Corresponding functions f_g and f_h are differ also. Indeed, for any $i \in e$ the set e is contained in a finite sequence $e_0(i) \ll \dots \ll e_{n_i}(i)$ of all sets that contain i . if its number in a such sequence is equal to k , then

$$f_g(i) = \text{span}\{\dots, g(e_k), \dots\} \neq \text{span}\{\dots, h(e_k), \dots\} = f_h(i).$$

Note that $e \in D$ and that $f_g(i) \neq f_h(i)$ for all $i \in e$.

Hence $f_g(i)$ and $f_h(i)$ generate different elements $\mathfrak{x}(f)$ and $\mathfrak{x}(h)$ of $(X_\alpha)_D$; moreover $\|\mathfrak{x}(f) - \mathfrak{x}(h)\| \geq \varepsilon$.

Since $\text{card}({}^G A) = \aleph^\tau$, and different elements $f, h \in {}^G A$ generate different elements $\mathfrak{x}(f)$ and $\mathfrak{x}(h)$ of $(X_\alpha)_D$ with $\|\mathfrak{x}(f) - \mathfrak{x}(h)\| \geq \varepsilon$, it is clear that $\dim((X_\alpha)_D) \geq \aleph^\tau$. $\square$

REMARK 3. *It may be proved that for any infinite-dimensional Banach space X and any countably incomplete ultrafilter D ,*

$$\dim((X)_D) = (\dim((X)_D))^\omega.$$

Hence, by using ultrapowers, cannot be obtained any space $(X)_D$, which dimension $\aleph$ is of countable cofinality (i.e. such that $\text{cf}(\aleph) = \omega$; e.g., $\aleph_\omega$; $\beth_{\omega_\omega}$ and so on).

THEOREM 9. *For any infinite dimensional Banach space X the corresponding class X^f contains spaces of every infinite cardinality (i.e., for any cardinal $\aleph \geq \omega$, $X^f \cap \mathcal{B}_\aleph \neq \emptyset$).*

PROOF. Let $\dim(X) = \tau$. If $\aleph = \omega$ then we choose a countable sequence $\{A_i : i < \omega\}$ of finite dimensional subspaces of X which is dense in a set $H(X)$ of all different finite dimensional subspace of X (isometric subspaces in $H(X)$ are identified), equipped with a metric topology, which is induced by the Banach-Mazur distance. Clearly, $X_0 = \text{span}\{A_i : i < \omega\} \hookrightarrow X$ is separable and is finite equivalent to X .

If $\aleph \leq \tau$ then as a representative of $X^f \cap \mathcal{B}_\aleph$ may be chosen any subspace of X of dimension τ that contains a subspace X_0 .

If $\tau < \aleph$ then, by the theorem 8 it may be found an ultrapower $(X)_D$ of dimension $\geq \aleph$. Certainly, $(X)_D \sim_f X$. Choose any subspace of $(X)_D$ of dimension $\aleph$, which contains a subspace $d_X X$. $\square$

7. Classes of ω -equivalence

If one replace in the definition of finite representability finite dimensional subspaces with infinite dimensional ones, it will be obtained a new relation of ω -representability.

DEFINITION 5. *Let X, Y be Banach spaces. X is said to be ω -representable in Y , shortly: $X <_\omega Y$, if for any separable subspace $E \hookrightarrow X$ there exists a separable $F \hookrightarrow Y$, which is isomorphic to E . X and Y are ω -equivalent, $X \sim_\omega Y$, if $X <_\omega Y$*

and $Y <_\omega X$. Any $X \in \mathcal{B}$ generates a class X^ω of all Banach spaces that are ω -equivalent to X :

$$X^\omega = \{Y \in \mathcal{B} : X \sim_\omega Y\}$$

The theorem 9 is not remain valid if one replaces X^f with X^ω . Of course, for some Banach spaces X a class X^ω also may contain spaces of arbitrary large dimension (e.g. if $X = l_p$ or $X = L_p$ for some $p \in [1, \infty]$).

It is easy to show that any Banach space X , which contains a subspace with a symmetric basis, generates a class X^ω , which contains spaces of arbitrary infinite dimension.

However, a method of ultrapowers cannot be used in a general case to construct other than X spaces from a class X^ω . Indeed, it is known (cf. [11]) that for any infinite dimensional Banach space X its ultrapower $(X)_D$ by a countably incomplete ultrafilter D is universal for all separable Banach spaces Y which are finitely representable in X . Hence, the following proposition is true.

PROPOSITION 1. *For any infinite dimensional Banach space and each pair of countably incomplete ultrafilters D and E spaces $(X)_D$ and $(X)_E$ are ω -equivalent.*

Another example gives a hereditarily indecomposable reflexive separable space X_{HI} , that was constructed in [2]. Obviously, a class $(X_{HI})^\omega$ cannot contain any non-separable space (since any non-separable reflexive Banach space is decomposable).

REMARK 4. *Let X be a Banach space of dimension $\kappa \geq v$, where v is the first measurable cardinal, and D be an ω -complete free ultrafilter. It may be proved that every subspace of $(X)_D$ of dimension, strictly less than v , is isometric to a subspace of X . This means that in a such case $X \sim_\omega (X)_D$.*

Since a dimension of $(X)_D$ (by choosing of an ω -complete free ultrafilter) may be made arbitrary large, it may be conclude that if a dimension $\dim(X) \geq v$, where v is the first measurable cardinal, then a class X^ω contains spaces of any dimension $\geq \aleph_1$.

The (too large) dimension v may be essentially lowered.

THEOREM 10. *For any Banach space X of dimension $\dim(X) \geq \aleph_{\omega_1}$ and any cardinal $\kappa \geq \aleph_{\omega_1}$, the corresponding class X^ω contains a space Y_κ of dimension κ .*

The proof of this theorem, which involves some model theory (and a series of additional definitions) is contained in an appendix to this article..

REMARK 5. *From this theorem follows immediately that if a cardinal $\kappa(\omega)$ exists then any Banach space X of dimension $\dim(X) = \aleph_{\omega_1}$ contains a subspace with an unconditional basis. Indeed, since $\aleph_{\omega_1} < \kappa(\omega)$, from the equality $\dim(X) = \aleph_{\omega_1}$ follows that the class X^ω contains a space Y of dimension $\kappa(\omega)$. This space contains a subspace W with an unconditional basis (see [4]). Since $Y <_\omega X$, X also has a subspace isomorphic to W .*

REMARK 6. *Note that Theorem 5 makes more clear the role of the cardinal $\aleph_{\omega_1}$.*

8. Appendix

8.1. Some model theory. To express the notion of a Banach space in the logic consider a *language* $\mathcal{L}$ - a set of symbols, which includes besides of logical symbols (prepositions $\&, \vee, \neg$, quantifiers $\forall, \exists$ and variables u, v, x_1, x_2 , etc) also non-logical primary symbols:

- A binary functional symbol $+$;
- A countable number of unary functional symbols $\{f_q : q \in \mathbb{Q}\}$;
- An unary predicate symbol B .

Any Banach space $\mathfrak{X} = \langle X, \|\cdot\| \rangle$ may be regarded as a model for all logical propositions (or formulae) that are satisfied in X . A set $|\mathfrak{X}| = X$ is called a support (or an absolute) of a model $\mathfrak{X}$, in which all non logical symbols of the language $\mathcal{L}$ are interpreted as follows:

- $+$ ^{$\mathfrak{X}$} is interpreted as the addition of vectors from X ;
- (f_q) ^{$\mathfrak{X}$} for a given $q \in \mathbb{Q}$ is interpreted as a multiplication of vectors from X by a rational scalar q
- B ^{$\mathfrak{X}$} is interpreted as the unit ball $B(X) = \{x \in X : \|x\| \leq 1\}$.

The language $\mathcal{L}$ was introduced by J. Stern [11], who used for examine the Banach space theory the *first order logic*. In this logic any quantifier acts only on elements of a supports of corresponding structures (e.g., a quantification over either all countable subsets of X or over all natural n 's is forbidden); only formulae that contains only finite strings of symbols are admissible. We shall used a less restrictive *infinitary logic* $\mathcal{L}_{\omega_1, \omega}$, which may be obtained from the first order logic by the addition of two new rules.

1. If Φ is a set of formulae of $\mathcal{L}_{\omega_1, \omega}$, such that $\text{card}(\Phi) \leq \omega$, then $\&\Phi$ also is a formula of $\mathcal{L}_{\omega_1, \omega}$;
2. If φ is a formula of $\mathcal{L}_{\omega_1, \omega}$ and V is a finite set of variables then $(\forall V)\varphi$ is a formula of $\mathcal{L}_{\omega_1, \omega}$.

In the logic $\mathcal{L}_{\omega_1, \omega}$ all axioms of Banach spaces can be expressed instead of the axiom of completeness (its formulation requires a quantification over a countable set of variables). For this reason (and for advantages of a countable language) any Banach space X will be regarded as a $\mathbb{Q}$ -vector normed space. Obviously, to any pair of $\mathbb{Q}$ -vector normed spaces $\mathfrak{X}, \mathfrak{Y}$, which are isomorphic in a model theoretic sense (i.e. that are such that there exists an one-to-one map $u : |\mathfrak{X}| \rightarrow |\mathfrak{Y}|$, which holds all functions and predicates of $\mathcal{L}$) corresponds (under their completion, if it needed) a pair of isometric Banach spaces. Thus, the infinitary logic $\mathcal{L}_{\omega_1, \omega}$ will be sufficient for our needs.

If φ is a formula of $\mathcal{L}_{\omega_1, \omega}$, which is satisfied in a Banach space X , we shall write $X \models \varphi$. Let

$$Th(X) = \{\varphi \in \mathcal{L}_{\omega_1, \omega} : X \models \varphi\}.$$

Banach spaces X and Y are said to be *elementary equivalent in the logic* $\mathcal{L}_{\omega_1, \omega}$, shortly, $X \equiv_{\omega_1, \omega} Y$, if $Th(X) = Th(Y)$.

The classical Scott's theorem asserts that for any countable system $\mathfrak{M}$ in a countable language for the logic $\mathcal{L}_{\omega_1, \omega}$ there exists such $\mathcal{L}_{\omega_1, \omega}$ -formula $\psi(\mathfrak{M})$ that the condition $\mathfrak{N} \models \psi(\mathfrak{M})$ for any other system $\mathfrak{N}$ implies that systems $\mathfrak{M}$ and $\mathfrak{N}$ are isomorphic. In our case, i.e., in the Banach space setting, this theorem means

that for any separable Banach space X the condition $X \equiv_{\omega_1, \omega} Y$ implies that X and Y are isometric.

As was noted by O.V. Belegadek (a personal communication), only small changes in the proof of the Scott's theorem are needed to do, to obtain the following its improvement:

For any system $\mathfrak{M}$ in a countable language for the logic $\mathfrak{L}_{\omega_1, \omega}$ there exists such $\mathfrak{L}_{\omega_1, \omega}$ -formula $\xi(\mathfrak{M})$ that a condition $\mathfrak{N} \models \psi(\mathfrak{M})$ for any other system $\mathfrak{N}$ implies that systems $\mathfrak{M}$ and $\mathfrak{N}$ have isomorphic countable subsystems.

From this immediately follows the following result.

THEOREM 11. *Let X, Y are Banach spaces. If $X \equiv_{\omega_1, \omega} Y$ then $X \sim_{\omega} Y$.*

PROOF. From the preceding improvement of the Scott's theorem follows that $X \sim_{\omega} Y$ if and only if $X \models Th(Y)_{\forall}$ and $Y \models Th(X)_{\forall}$, where for any theory T (i.e. for any set of $\mathfrak{L}_{\omega_1, \omega}$ -formulae) $T_{\forall}$ denotes a set of all its universal consequences. Of course, $T_{\forall} \subseteq T$ for any theory T . Therefore $X \equiv_{\omega_1, \omega} Y$ implies that $X \sim_{\omega} Y$. $\square$

The next notion that will be needed is a notion of a *Hanf number* of the logic $\mathfrak{L}_{\omega_1, \omega}$.

Let X be a $\mathbb{Q}$ -vector normed space. Let $ModTh(X) = \{Y : Y \equiv_{\omega_1, \omega} X\}$, where Y is a $\mathbb{Q}$ -vector normed space too. The Hanf number of the logic $\mathfrak{L}_{\omega_1, \omega}$ is the least cardinal $h(\mathfrak{L}_{\omega_1, \omega})$ with the property:

If a model $\mathfrak{M}$ for this logic is of cardinality $card|\mathfrak{M}| \geq h(\mathfrak{L}_{\omega_1, \omega})$, then for any cardinal $\kappa \geq h(\mathfrak{L}_{\omega_1, \omega})$ there exists a model $\mathfrak{N} \equiv_{\omega_1, \omega} \mathfrak{M}$ of cardinality $card|\mathfrak{N}| = \kappa$.

In our case this means that for each Banach space X of dimension $dim(X) \geq h(\mathfrak{L}_{\omega_1, \omega})$ and for any $\kappa \geq h(\mathfrak{L}_{\omega_1, \omega})$ there exists a Banach space $Y \equiv_{\omega_1, \omega} X$ of dimension $dim(Y) = \kappa$.

The exact meaning of the Hanf number for the logic $\mathfrak{L}_{\omega_1, \omega}$ is known (see [13]):

$$h(\mathfrak{L}_{\omega_1, \omega}) = \mathfrak{J}_{\omega_1}$$

Now we are ready to present a proof of the Theorem 7.

8.2. Proof of the theorem 10.

PROOF. Let X be a Banach space of dimension $dim(X) = \mathfrak{J}_{\omega_1}$. Consider X to be a model for the infinitary logic $\mathfrak{L}_{\omega_1, \omega}$. Since its dimension is equal to the Hanf number $h(\mathfrak{L}_{\omega_1, \omega})$, for any $\kappa \geq \mathfrak{J}_{\omega_1}$, then there exists $Y \equiv_{\omega_1, \omega} X$ of dimension $dim(Y) = \kappa$. Clearly, the condition $Y \equiv_{\omega_1, \omega} X$ implies that $Y \sim_{\omega} X$. The last equivalence proves the theorem. $\square$

REMARK 7. *Let X be a Banach space of dimension $\kappa > \omega_1$. Then there exists a Banach space $Y \equiv_{\omega_1, \omega} X$ of dimension $dim(Y) = \omega_1$. This follows from the general result of R.L. Vaught [13]: the Löwenheim number $l(\mathfrak{L}_{\omega_1, \omega})$ for a countable language $\mathfrak{L}$ (i.e. the least cardinal τ such that every $\mathfrak{L}$ -model $\mathfrak{A}$ of cardinality $\varsigma > \tau$ has a subsystem $\mathfrak{B} \models Th(\mathfrak{A})$ of cardinality τ) is equal to ω_1 .*

ACKNOWLEDGEMENT 1. *Author wish to express his thanks to O.V. Belegadek for helpful advises and recommendations*

9. References

1. Lindenstrauss J. *The geometric theory of the classical Banach spaces*, Actes du Congres Intern. Math. Nice 1970, v. **2** (1971) 365-372
2. Gowers W.T., Maurey B. *The unconditional basic sequence problem*, Journal AMS **6** (1993) 851-874
3. Ketonen J. *Banach spaces and large cardinals*, Fund Math. **81** (1974) 291-303
4. Tokarev E.V. *Notion of indiscernibleness and on the p -Mazur property in Banach spaces* (transl. from Russian) Ukrainian Mathematical Journal **37** (1985) p.180-184
5. Tokarev E.V. *On one P. Terenzi's question*, Deposited in VINITI no.**325-B88** (1988) 1-7 (in Russian)
6. Terenzi P. *On the Banach spaces of large density character*, Riv. Mat. Univ. Parma **12:4** (1986) 111-120
7. Erdős P., Rado R. *A partition calculus in set theory*, Bull. AMS **62** (1956) 427-489
8. Brunel A., Sucheston L. *On J - convexity and some ergodic super properties of Banach spaces*, Trans. AMS **204** (1975) 79-90
9. Heinrich S. *Ultraproducts in Banach space theory*, J. Reine Angew. Math. **313** (1980) 72-104
10. Chang C. C., Keisler H. J. *Model Theory*, Amsterdam: North-Holland 1973
11. Stern J. *The problem of envelopes for Banach spaces*, Israel J. Math. **24:1** (1976) 1-15
12. Hanf W. *Incompactness in languages with infinitely long expressions*, Fund. Math. **13** (1964) 309-324
13. Vaught R.L. *Denumerable models of complete theories*, in: Infinitistic Methods; Pergamon Press (1961) 303-321

B.E. UKRECOLAN, 33-81, ISKRINSKAYA STR.,61005, KHARKIV-5, UKRAINE
 E-mail address: evtokarev@yahoo.com